



HiFluoro Pseudomonas HiVeg Agar Base

MV1469

HiFluoro Pseudomonas HiVeg Agar Base is recommended for selective isolation of *Pseudomonas aeruginosa* from clinical and non-clinical specimens by fluorogenic method.

Composition**

Ingredients	Gms / Litre
HiVeg Peptone No.2	18.000
Magnesium chloride	1.400
Potassium sulphate	10.000
Cetrimide	0.300
Fluorogenic mixture	2.050
Agar	15.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 46.75 grams in 1000 ml distilled water containing 10ml glycerol. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Pseudomonas aeruginosa (also known as *Pseudomonas pyocyanea*) is a gram-negative, aerobic, rod-shaped bacterium. Like other Pseudomonads, *P. aeruginosa* secretes a variety of pigments, including pyocyanin (blue-green), fluorescein (yellow-green and fluorescent), and pyorubin (red-brown). King et al developed Pseudomonas Agar P (i.e. King A media) for enhancing pyocyanin and pyorubin production and Pseudomonas Agar F (i.e. King B media) for enhancing fluorescein production (1). HiFluoro Pseudomonas HiVeg Agar Base is prepared by completely replacing animal based peptones with veg peptones. This media is a modification of HiFluoro Pseudomonas Agar Base which is devised based on the formula described by King et al. (1) except fluorogenic mixture. It is used as the selective medium for the isolation of *P. aeruginosa* from pus, sputum and drains etc.

Cetrimide (Cetyltrimethylammonium bromide) is incorporated in the medium to inhibit bacteria other than *P. aeruginosa*. It acts as a quaternary ammonium compound, cationic detergent that causes nitrogen and phosphorus to be released from bacterial cells other than *P.aeruginosa*. *P.aeruginosa* cleaves the fluorogenic compound to release the fluorogen which produces a visible fluorescence under long wave UV light.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber colourec opalescent gel forms in Petri plates

Reaction

Reaction of 4.675 % w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Fluorescence (under uv)
Cultural Response				
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	>=50%	positive
<i>Stenotrophomonas maltophilia</i> ATCC 13637	>=10 ³	inhibited	0%	
<i>Staphylococcus aureus</i> ATCC 25923	>=10 ³	inhibited	0%	
<i>Escherichia coli</i> ATCC 25922	>=10 ³	inhibited	0%	

Storage and Shelf Life

Store dehydrated and prepared medium at 2-8°C in tightly closed container. Use before expiry period on the label.

Reference

1.King, Ward and Raney, 1954, J. Lab. Clin. Med., 44:301.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.